



BIOGEME: a free package for the estimation of discrete choice models

Michel Bierlaire, EPFL

Conference paper STRC 2003

STRC 3rd Swiss Transport Research Conference
Monte Verita, Ascona, March 2003

BIOGEME: a free package for the estimation of discrete choice models

Michel Bierlaire, Dpt of Mathematics, EPFL, Lausanne

Phone: +41.21.693.25.37

Fax: +41.21.693.55.70

Email: michel.bierlaire@epfl.ch

Abstract

Keywords: logit model, cross-nested model, GEV model, random utility, transportation demand

Swiss Transport Research Conference – STRC 2003 – Monte Verita

1 Introduction

Discrete choice models in general, and random utility models in particular, have been extensively used in several fields of applications for the last three decades. The theoretical derivation of these models is well documented in the literature (Luce, 1959, McFadden, 1981, Ben-Akiva and Lerman, 1985, Anderson et al., 1992, Hensher and Johnson, 1981, Horowitz et al., 1986, Bierlaire, 1998, Ben-Akiva and Bierlaire, 1999). Recently, new sophisticated models have been proposed in the literature. Among them, two main categories can be identified: Generalized Extreme Value models, and mixed logit models.

Generalized Extreme Value (GEV) models have been proposed 25 years ago by McFadden (1978). It is actually a family of models, consistent with random utility theory. Since then, only a few members have been exploited, the Multinomial Logit model and the Nested Logit model being the most popular (Ben-Akiva and Lerman, 1985). Recently, research on the Cross-Nested logit model (Small, 1987, Vovsha, 1997, Vovsha and Bekhor, 1998, Ben-Akiva and Bierlaire, 1999, Papola, 2000, Bierlaire, 2001, Wen and Koppelman, 2001, Swait, 2001) has extended the number of GEV models used in practice. Also, Daly (2001) and Bierlaire (2002) have proposed new theoretical results providing an operational representation of GEV models.

Mixed or hybrid logit models (Ben-Akiva and Bolduc, 1996, McFadden and Train, 1997, Bhat, 2001, Ben-Akiva et al., 2002) combine different model structures into a richer framework. A Mixed GEV model can be roughly described as a GEV model containing random parameters, which are normally distributed.

The level of sophistication of these models enables to capture a wide range of situations. The price to pay is their rather complicated formulation, and the lack of appropriate estimation procedure. For example, Vovsha (1997) proposes a heuristic procedure for the estimation of Cross-Nested Logit models, which appears not to be valid.

Biogeme (**B**ierlaire's **O**ptimization package for **GEV** **M**odels **E**stimation) is an open source package designed to estimate a wide variety of random utility models, based on state-of-the-art optimization algorithms. The motivation for developing Biogeme is to provide researchers with an appropriate and efficient tool enabling to explore new models, focusing on their specification without worrying about the estimation part. In this paper, we describe the general design of the Biogeme package, and we illustrate the capabilities of Version 0.6.

2 Random utility models

A random utility model is designed to forecast the choice of an individual n among a finite and discrete set of alternatives C_n . The main assumption is that each individual associates a quantity, called *utility*, to each alternative in C_n , and selects the alternative with the highest utility. The utility associated by individual n to alternative i , denoted by U_{in} is a random variable such that

$$U_{in} = V_{in} + \varepsilon_{in} \quad (1)$$

where $V_{in} \in \mathbb{R}$ is the deterministic, or systematic, component of the utility, and ε_{in} is a random term. If z_{in} is a vector of attributes of alternative i for individuals n , and S_n is a vector of socio-economical characteristics for individual n , we have

$$V_{in} = V_{in}(\beta, z_{in}, S_n), \quad (2)$$

where β is a vector of unknown parameters to be estimated. For simplification, it is common practice to merge z_{in} and S_n into a vector of attributes, denoted by x_{in} . Therefore, we have a simpler formulation

$$V_{in} = V_{in}(\beta, x_{in}) \quad (3)$$

The probability that individual n selects alternative i is given by

$$P(i|C_n) = P(U_{in} \geq U_{in} \quad \forall j \in C_N). \quad (4)$$

In order to obtain an operational choice model, specific assumptions must be made about the functional form of V_{in} and the distribution of ε_{in} . We briefly review the assumptions which are relevant for the Biogeme package.

The most common functional form adopted for V_{in} is a linear-in-parameters definition, that is

$$V_{in} = \sum_j \beta_j x_{inj}. \quad (5)$$

Less common in the literature, nonlinear formulations may also be used. The Box-Tukey transform of attributes is a typical example, that is

$$V_{in} = \sum_j \beta_j \frac{(x_{inj} + \alpha_{inj})^{\lambda_{inj}} - 1}{\lambda_{inj}}. \quad (6)$$

where $x_{inj} + \alpha_{inj}$ must be non negative, α_{inj} and λ_{inj} are unknown parameters to be estimated. Box-Tukey transforms allow to capture a wide range of

non-linearities. Indeed, in addition to the obvious exponential transformation of the attributes, linear ($\lambda_{inj} = 1$) and logarithm ($\lambda_{inj} = 0$) transformations are interesting special cases.

Before discussing the assumptions about the distribution of ε_{in} , we analyze the mean and the variance of the random variable. The mean can be considered as a specific parameter of the utility function (called *Alternative Specific Constant*), capturing a bias toward that alternative. In that case, ε_{in} is decomposed into

$$\varepsilon_{in} = \beta_i^0 + \tilde{\varepsilon}_{in} \quad (7)$$

so that the expectation of $\tilde{\varepsilon}_{in}$ can be set to any arbitrary value, typically zero. Therefore, if an alternative specific constant is included in the utility function, the mean can be assumed to be zero without loss of generality.

The case of the variance is discussed by first noting that

$$P(U_{in} \geq U_{in}) = P(\nu U_{in} \geq \nu U_{in}) \quad \forall \nu > 0.$$

Therefore, using $U_{in} = V_{in} + \varepsilon_{in}$ or $\nu U_{in} = \nu V_{in} + \nu \varepsilon_{in}$ yields to the exact same probability model. As

$$\text{Var}(\nu U_{in}) = \nu^2 \text{Var}(U_{in}),$$

the choice of ν determines the variance. Therefore, the variance of the random parameter is directly linked to its scale, and can be arbitrarily imposed. $\nu = 1$ is a typical choice.

The above discussion is valid only for a given individual. It is important to realize that assuming constant mean and variance over the population may be a strong and unrealistic assumption. In most applications, those quantities are different for various groups of the population. The variations of the mean across the population are captured by dummy parameters associated with socio-economic characteristics. Variations of the variance (or, equivalently, of the scale parameter ν) are more complicated to capture, as they introduce a nonlinearity in the utility function. If we consider a heterogeneous population composed of identified groups, the utility of each group is scaled by a different factor. In that case, if individual n belongs to group g_n , we generalize (5) as

$$V_{in} = \nu_{g_n} \sum_j \beta_j x_{inj}. \quad (8)$$

and (6) as

$$V_{in} = \nu_{g_n} \sum_j \beta_j \frac{(x_{inj} + \alpha_{inj})^{\lambda_{inj}} - 1}{\lambda_{inj}}, \quad (9)$$

where ν_{g_n} is the unknown scale parameter associated with group g_n , to be estimated. A typical example is when Revealed Preference (RP) data are combined with Stated Preference (SP) data (Ben-Akiva et al., 1994, Ben-Akiva and Morikawa, 1990). Note that (5), (6) and (8) are special cases of (9), where some parameters are set to fixed values. Whatever the assumptions about the distribution of ε_{in} , Biogeme allows to estimate parameters β , λ , α and ν in (9) from the data, if they are identifiable.

In the literature, there are typically two families of models, based on two types of assumptions about the error term ε_{in} . The *Probit* family assumes that ε_{in} captures the sum of many independent sources of errors, and invokes the central-limit theorem to assume that ε_{in} follows a normal distribution. The *Generalized Extreme Value* (GEV) family, assumes that ε_{in} captures the largest of many independent sources of errors and, consequently, has an Extreme Value distribution (Gumbel, 1958).

The Generalized Extreme Value (GEV) model has been derived from the random utility paradigm by McFadden (1978). This general model consists of a large family of models that include the Multinomial Logit, the Nested Logit and the Cross-Nested Logit models. The probability of choosing alternative i within the choice set C of a given choice maker is

$$P(i|C) = \frac{y_i \frac{\partial G}{\partial y_i}(y_1, \dots, y_J)}{\mu G(y_1, \dots, y_J)} \quad (10)$$

where J is the number of available alternatives, $y_i = e^{V_i}$, V_i is the deterministic part of the utility function associated to alternative i , and G is a μ -GEV function. A μ -GEV function is a differentiable function defined on \mathbb{R}_+^J with the following properties:

1. $G(y) \geq 0$ for all $y \in \mathbb{R}_+^J$,
2. G is homogeneous of degree $\mu > 0$, that is $G(\lambda y) = \lambda^\mu G(y)$, for $\lambda > 0$,
3. $\lim_{y_i \rightarrow +\infty} G(y_1, \dots, y_i, \dots, y_J) = +\infty$, for each $i = 1, \dots, J$,
4. the k th partial derivative with respect to k distinct y_i is non-negative if k is odd and non-positive if k is even that is, for any distinct indices $i_1, \dots, i_k \in \{1, \dots, J\}$, we have

$$(-1)^k \frac{\partial^k G}{\partial x_{i_1} \dots \partial x_{i_k}}(x) \leq 0, \quad \forall x \in \mathbb{R}_+^J. \quad (11)$$

Although this condition is never stated in the literature, it is also required that $G(x) \neq 0$.

The homogeneity of G and Euler's theorem give

$$P(i|C) = \frac{e^{V_i + \ln G_i(\dots)}}{\sum_{j=1}^J e^{V_j + \ln G_j(\dots)}}, \quad (12)$$

where $G_i = \frac{\partial G}{\partial y_i}$. This is equivalent to assume that the joint distribution of the error terms $\varepsilon_1, \dots, \varepsilon_J$ is

$$F(\varepsilon_1, \dots, \varepsilon_J) = e^{-G(e^{-\varepsilon_1}, \dots, e^{-\varepsilon_J})} \quad (13)$$

The Multinomial Logit (MNL) model is a member of this model family with

$$G(y) = \sum_{i=1}^J y_i^\mu. \quad (14)$$

where $\mu > 0$. The Nested Logit (NL) model is also a member of the GEV family. Contrarily to the MNL, the NL includes $M + 1$ parameters, where M is the number of nests. The generating function is

$$G(y; \mu_1, \dots, \mu_M) = \sum_{m=1}^M \left(\sum_{i \in C_m} y_i^{\mu_m} \right)^{\frac{\mu}{\mu_m}} \quad (15)$$

where C_m is the set of alternatives belonging to nest m . It complies with the GEV conditions if $\mu_m \geq \mu > 0$, for all m .

The Cross-Nested Logit (CNL) model is also a member of the GEV family. Several formulations have been proposed in the literature (Small, 1987, Vovsha, 1997, Ben-Akiva and Bierlaire, 1999, Papola, 2000, Wen and Koppelman, 2001). They are all based on the same formulation, and vary with regard to the parameters that are kept fixed. A detailed analysis of this model, including the elasticities, have been recently proposed by Wen and Koppelman (2001). However, they call it the *generalized nested logit*. The generating function proposed by Ben-Akiva and Bierlaire (1999), with $M(J + 1) + 1$ parameters, where M is the number of nests, is given by

$$G \left(y; \mu_1, \dots, \mu_M, (\alpha_{jm})_{j=1, \dots, J}^{m=1, \dots, M} \right) = \sum_{m=1}^M \left(\sum_{j=1}^J \alpha_{jm} y_j^{\mu_m} \right)^{\frac{\mu}{\mu_m}}. \quad (16)$$

Bierlaire (2001) has shown that the following conditions are sufficient for (16) to comply with the GEV conditions:

1. $\mu_m \geq \mu > 0, m = 1, \dots, M,$
2. $\alpha_{jm} \geq 0, j = 1, \dots, J, m = 1, \dots, M,$
3. $\sum_m \alpha_{jm} > 0, j = 1, \dots, J.$

The Network GEV (NGEV) model, proposed by Bierlaire (2002), and based on a model formulation by Daly (2001), is a general representation of GEV models, based on a network structure. A parameter is associated with each node and each arc of this network. We refer the reader to Bierlaire (2002) for more details.

Thanks to Denis Bolduc, Biogeme v0.6 is also able to estimate Logit Kernel models with error components. A Logit-Kernel model is such that the utility function of an alternative is

$$U_i = V_i + \sum_{j=1}^p \sigma_j \nu_j + \varepsilon_i \quad (17)$$

where σ_j are unknown parameters to be estimated, and ν_j are normal random variables $N(0, 1)$.

In the rest of the paper, we denote by β the unknown parameters associated with the utility function (that is, parameters β, λ, α and ν in (9)), and by γ the unknown parameters associated with a specific GEV model (that is, parameters μ_m in the NL, CNL and NGEV models, and parameters α_{im} in the CNL and NGEV models).

3 Maximum likelihood estimation

The estimation of unknown parameters by maximum likelihood is a standard technique. An observation k consists in a set of values for the set of attributes x_{in} , denoted x_{in}^k , and an observed choice. The attributes are associated both with the individual n and the alternative i . The probability for the model to reproduce the observed choice is given by $P_{in}^k(\beta, \gamma) = P_{in}(\beta, \gamma, x_{in}^k)$, where P_{in} is the probability function corresponding to the model under consideration (like (10) for the GEV model). If a sample of K observations is available, the probability for the model to reproduce the whole sample is called the *likelihood*, and is given by

$$\mathcal{L}^*(\beta, \gamma) = \prod_{k=1}^K P_{in}^k(\beta, \gamma). \quad (18)$$

The maximum likelihood estimators $\hat{\beta}$ and $\hat{\gamma}$ are given by

$$(\hat{\beta}, \hat{\gamma}) = \operatorname{argmax}_{\beta, \gamma} \mathcal{L}(\beta, \gamma), \quad (19)$$

where

$$\mathcal{L}(\beta, \gamma) = \ln \mathcal{L}^*(\beta, \gamma) = \sum_{k=1}^K \ln P_{in}^k(\beta, \gamma) \quad (20)$$

is the log-likelihood function. In some cases, the observations are weighted in order to adjust their relative importance in the sample according to their relative importance in the population. In that case, a weight ω_k is associated with each observation, and the log-likelihood function is then

$$\mathcal{L}(\beta, \gamma) = \sum_{k=1}^K \omega_k \ln P_{in}^k(\beta, \gamma) \quad (21)$$

Problem (19) is a nonlinear programming problem, usually non concave. Moreover, it is sometimes necessary (and most of the time useful) to impose constraints on β and γ . For example, the condition $\mu_m \geq \mu$ is necessary for the validity of the NL and CNL models. Also, normalisation conditions may be imposed on the parameters. Note that Biogeme allows to impose bound constraints, linear equality and inequality constraints, and nonlinear equality constraints on the parameters.

Biogeme contains three different optimization algorithms. CFSQP is a C implementation of the FSQP optimization algorithm developed by E.R. Panier, A.L. Tits, J.L. Zhou, and C.T. Lawrence (see Lawrence et al., 1997). SolvOpt (Solver for local optimization problems) by Kuntsevich and Kappel (1997) implements a version of a minimization method with space dilation by Shor (1985). And DONLP2 is a sequential equality constrained quadratic programming method, developed by Spellucci (n.d.). The algorithm is described by Spellucci (1998a) and Spellucci (1998b).

All those algorithms identify a local optimum of (19). In the (rare) cases where the objective function is concave, and the constraints are convex, the local optimum is also global.

3.1 Comparison

We provide in Table 1 a comparison of the final log-likelihood for the models available on the website, and the time it took to estimate them with CFSQP,

DONLP2 and SOLVOPT on a Dell Inspiron 8200 running Linux RedHat 7.3. Entries with **** correspond to failure of convergence of the algorithm. The examples are describe in Bierlaire (2003). We provide here some comments about the results.

- CFSQP is most of the time the fastest algorithm, followed by DONLP2 and finally by SOLVOPT, the slowest of the three.
- Biogeme may be significantly slower for general NGEV, as it does not exploit the special structure of the model.
- On the difficult problem number 10, CFSQP was much slower than DONLP2 with the CNL version (EX10), and did not even converge after 1000 iterations on the NGEV version (NGEV10).
- For Logit Kernel models (examples EX13 to EX18) the final log-likelihood may vary from one estimation to the next, as the normally distributed random variables are simulated based on a Monte-Carlo procedure.

4 Biogeme Packages

BIOGEME is a freeware designed for the development of research in the context of discrete choice models in general, and of GEV models in particular. All information relative to BIOGEME is maintained at

`http://roso.epfl.ch/biogeme`

where a detailed tutorial is available (Bierlaire, 2003). We cite here the main features of the packages.

BIOGEME has been developed on Linux, but a Windows version is available. With the distribution of Biogeme, there are two additional utilities. Bioroute helps preparing the input files for Biogeme in the context of a route choice analysis, and Biosim is designed to perform simulations with a given model. Biogeme is invoked by the following command

```
biogeme model_name sample_file
```

If the name of the model is `mymodel`, say, Biogeme reads the following files:

- a file containing the parameters controlling the behavior of Biogeme: `mymodel.par`,
- a file containing the model specification: `mymodel.mod`,
- a file containing the data: `sample.dat`,

and generates the following output files:

- a file reporting the results of the estimation: `mymodel.rep`,
- the same file in HTML format: `mymodel.html`,
- a file containing the specification of the estimated model, in the same format as the model specification file: `mymodel.res`,
- a file containing some statistics on the data: `mymodel.sta`.

For most users, the parameter file is edited only to select a specific optimization algorithm. The data file contains in its first line a list of labels corresponding to the available data, and that each subsequent line contains the exact same number of numerical data, each row corresponding to an observation. The model specification file is based on a syntax designed to define a wide range of models, with several sections. We enumerate here the most important sections.

Section [Beta] Each line of this section corresponds to a parameter β in (9).

Five entries must be provided for each parameter: its name, a default value, a lower bound and an upper bounds on the valid values and a binary status, specifying if the value of the parameter must be estimated or kept at its default value. Note that this section is independent of the specific model to be estimated, as it captures only the deterministic part of the utility function. Here is an example.

```
[Beta]
// Name Value          LowerBound   UpperBound   status
ASC1   -5.22e-02           -1.0         1.0         0
ASC2         0.0              -1.0         1.0         1
ASC3   -4.06e-01           -1.0         1.0         0
ASC4         0.0              -1.0         1.0         1
BETA1  -2.06e-02           -1.0         1.0         0
BETA2  -2.19e-02           -1.0         1.0         0
```

Section [Utilities] For each alternative in the model, the following information must be provided in this section: the numerical identifier of the alternative, the name of the alternative, the availability condition and the linear-in-parameter utility function. For example,

```
[Utilities]
//Id Name Avail linear-in-parameter expression
100 Alt1 avail1 BETA_COST * COST1 + BETA_TIME * TIME1 + ASC1 * one
200 Alt2 avail2 BETA_COST * COST2 + BETA_TIME * TIME2 + ASC2 * one
300 Alt3 avail3 BETA_COST * COST3 + BETA_TIME * TIME3 + ASC3 * one
400 Alt4 avail4 BETA_COST * COST4 + BETA_TIME * TIME4
```

where lines starting by // are ignored by Biogeme and used for comments.

Section [Box-Cox] Each line of this section corresponds to a parameter λ in (9). The following entries must be provided: the name of the attribute x_{inj} , a default value, a lower bound and an upper bounds on the valid values and a binary status, specifying if the value of the parameter must be estimated or kept at its default value.

Section [Box-Tukey] Each line of this section corresponds to a parameter α in (9). The following entries must be provided: the name of the attribute x_{inj} , a default value, a lower bound and an upper bounds on the valid values and a binary status, specifying if the value of the parameter must be estimated or kept at its default value.

Section [Expressions] In this section are defined all expressions appearing either in the availability conditions or in the utility functions of the alternatives. If the expression is readily available from the data file, it can be omitted in the list. It is especially useful to tests alternative model specifications (like nonlinear effects) without modifying the data file.

Section [Choice] The user provides here the formula to compute the identifier of the chosen alternative in the data file. Typically, a “choice” entry will be available directly in the file, but any formula can be used to compute it.

Section [Weight] The user provides here the formula to compute the weights ω_k in (21). Ideally, the sum of the weights should be equal to the total number of observations, although it is not required.

Section [Group] The user provides here the formula to compute the group ID of the observed individual. Typically, a “group” entry will be available directly from the data file, but any formula can be used to compute it. A different scale parameter μ_{g_n} will be estimated for each group.

Section [Scale] Each line of this section corresponds to a parameter ν_{g_n} in (9), one per group. The following entries must be provided: the group number, a default value, a lower bound and an upper bound on the valid values and a binary status, specifying if the value of the parameter must be estimated or kept at its default value.

Section [Model] Selects the GEV model. Valid entries are **\$MNL** for Multinomial Logit model, **\$NL** for single level Nested Logit model, **\$CNL** for Cross-Nested Logit model (in the sense described in Ben-Akiva and Bierlaire (1999)), **\$NGEV** for Network GEV model, and **\$LK** for Logit Kernel model.

Section [NLNests] This section is relevant only if the **\$NL** option has been selected. Each row of this section corresponds to a nest. The following entries are required: the nest name, a default value, a lower bound and an upper bound on the valid values of the nest parameter μ_m , and a binary status, specifying if the value of the parameter must be estimated or kept at its default value. And finally, the list of alternatives belonging to the nest. Similar sections must be defined if a CNL or a NGEV model are to be estimated.

Section [ConstraintNestCoef] In this section, the user can constraint nests parameters to be equal, with the following syntax.

```
NEST_A = NEST_B
```

Section [ConstantProduct] In this section, the user may constrain the product of two coefficients to a given value. The syntax for the constraint $\beta_1\beta_2 = 3.1415$ is

```
[ConstantProduct]
BETA1    BETA2    3.1415
```

Section [Ratios] The user defines here the ratio of parameters that must be computed. Typically, the value of time is the ratio for the time parameter by the cost parameter.

Section [LinearConstraints] In this section, the user can define a list of linear constraints, in one of the following syntaxes:

1. Formula = number,
2. Formula \leq number,
3. Formula \geq number.

For example, the constraint

$$\sum_i ASC_i = 0.0$$

is written

$$ASC1 + ASC2 + ASC3 + ASC4 + ASC5 + ASC6 = 0.0$$

and the constraint

$$\mu \leq \mu_j$$

is written

$$MU - MUJ \leq 0.0$$

or

$$MUJ - MU \geq 0.0$$

Section [NonLinearEqualityConstraints] In this section, the user can define a list of nonlinear equality constraints of the form

$$h(x) = 0.0.$$

The section must contain a list of functions $h(x)$. For example, the constraint

$$\alpha_{a1}^{\mu_a} + \alpha_{b1}^{\mu_b} = 1$$

is written

```
[NonLinearEqualityConstraints]
ALPHA_A1 ^ MU_A + ALPHA_B1 ^ MU_B - 1.0
```

Section [LogitKernelSigmas] Each line of this section corresponds to a σ parameter associated with the normal terms in the utility functions for the Logit Kernel (17). Note that the sign of the parameter is meaningless, so it is good practice to specify symmetric bounds (that is, lower bound is the opposite of the upper bound).

Section [LogitKernelFactors] This section defines the factors of the Logit Kernel model. For each factor, the name of the associated parameter and the ID of the associated alternative must be specified.

The report file (`mymodel.rep`) contains the results of the maximum likelihood estimation of the model.

- The estimated value of the β parameters, with the associated standard error and the t -test. A star (*) is appended if the t -test fails, according to a threshold specified by the use in the parameter file (default threshold: 1.96).
- The estimated value of the μ parameter, with the associated standard error and the t -test.
- The estimated value of the GEV model parameters, with the associated standard error and the t -test. Note that the t -test is computed to compare the estimated value both to 0 and 1.
- The estimated value of the scale parameters, with the associated standard error and the t -test. Note that the t -test is computed to compare the estimated value to 1.
- A covariance/correlation analysis of pairs of estimated β parameters, sorted according to the t -test value.

A list of examples is available from the BIOGEME webpage, and are commented in Bierlaire (2003).

The package Biosim is invoked exactly like Biogeme, with the exact same input file. But instead of performing a parameter estimation, it uses the default value for each parameter, performs a sample enumeration and produces a Gnuplot file allowing for a graphical display of the model sensitivity. The output file `mymodel.enu` contains the result of the sample enumeration. For each observation in the sample, the following results are provided:




Figure 1: Example of a Gnuplot output

1. The choice actually reported in the sample file;
2. The probability given by the model for the chosen alternative;
3. For each alternative, the probability given by the model;
4. A list of simulated choice, based on random draws using the model.

The file `mymodel.gp` is an input file for the Gnuplot. It allows to graphically analyze the sensitivity of the model to modifications of one attribute (see, for instance, Figure 1).

Finally, the package `Bioroute` is used in the context of route choice analysis. Indeed, the explicit enumeration of paths may be cumbersome when specifying route choice models. Moreover, computing the size of the paths for Path-Size models (Ben-Akiva and Bierlaire, 1999, Ramming, 2001) or the cross-nested coefficients for the link-nested logit model (Vovsha and Bekhor, 1998) is tedious and subject to errors. The utility `BioRoute` is designed to help the analyst. `Bioroute` takes as input a full description of the network, and prepares the files needed by

Biogeme, that is the model specification file and the sample file. After Bioroute has been used, it is important to edit the generated specification file.

5 Future developments

Biogeme is in continuous development. Biogeme 0.7 will be able to estimate general nonlinear utility functions, and *Mixed GEV models*. Mixed GEV models are a generalization of Mixed Logit models, also called Hybrid Logit or Logit Kernel models (see McFadden and Train, 1997, Ben-Akiva et al., 2001, Bhat, 2001, Ben-Akiva et al., 2002). This is a major step that will allow a vast variety of models to be estimated, including the heteroscedastic model, the factor analytic with fixed loadings, the error component formulation and the general autoregressive process (see Ben-Akiva and Bierlaire, 1999, Walker, 2001 and Ben-Akiva et al., 2001). All these efforts are motivated by the same objective: provide researchers and practitioners with flexible tools to investigate a wide range of discrete choice models.

6 Appendix: derivatives

In addition to the programming burden, a major effort in developing packages for model estimation is the computation of the derivatives required by the optimization packages. We provide in this appendix the derivatives used in Biogeme for log-likelihood estimation of Multinomial, Nested and Cross-Nested logit models.

The log-likelihood function is defined by (21). The derivatives are trivially defined as

$$\frac{\partial \mathcal{L}}{\partial \beta} = \sum_{k=1}^K \frac{\partial \ln P_{in}^k(\beta, \gamma)}{\partial \beta} \quad (22)$$

and

$$\frac{\partial \mathcal{L}}{\partial \gamma} = \sum_{k=1}^K \frac{\partial \ln P_{in}^k(\beta, \gamma)}{\partial \gamma}. \quad (23)$$

Denoting by i the chosen alternative, V the J utilities, γ the ℓ model parameters, and ν the scale parameter, we have

$$P(i, V_1, \dots, V_J, \gamma_1, \dots, \gamma_\ell, \nu, \mu) = \frac{e^{\nu V_i + \ln G_i(e^{\nu V_1}, \dots, e^{\nu V_J}, \gamma_1, \dots, \gamma_\ell, \mu)}}{\sum_j e^{\nu V_j + \ln G_j(e^{\nu V_1}, \dots, e^{\nu V_J}, \gamma_1, \dots, \gamma_\ell, \mu)}}$$

and

$$\begin{aligned}\ln P(i, V_1, \dots, V_J, \gamma_1, \dots, \gamma_\ell, \nu, \mu) &= \nu V_i \\ &+ \ln G_i(e^{\nu V_1}, \dots, e^{\nu V_J}, \gamma_1, \dots, \gamma_\ell, \mu) \\ &- \ln \left(\sum_j e^{\nu V_j} G_j(e^{\nu V_1}, \dots, e^{\nu V_J}, \gamma_1, \dots, \gamma_\ell, \mu) \right).\end{aligned}$$

The derivatives with respect to β_k are given by

$$\frac{\partial}{\partial \beta_k} \ln P = \nu \frac{\partial V_i}{\partial \beta_k} + \frac{1}{G_i} \sum_{j=1}^J \frac{\partial G_i}{\partial x_j} e^{\nu V_j} \nu \frac{\partial V_j}{\partial \beta_k} - \frac{1}{\Delta} \sum_j e^{\nu V_j} \left(\nu \frac{\partial V_j}{\partial \beta_k} G_j + \sum_{n=1}^J \frac{\partial G_j}{\partial x_n} e^{\nu V_n} \nu \frac{\partial V_n}{\partial \beta_k} \right)$$

where

$$\Delta = \sum_j e^{\nu V_j} G_j.$$

Note that we do not assume here that the V_j are linear-in-parameters, so that $\partial V_j / \partial \beta_k$ may be not trivial.

The derivatives with respect to the model parameters γ_k are given by

$$\frac{\partial}{\partial \gamma_k} \ln P = \frac{1}{G_i} \frac{\partial G_i}{\partial \gamma_k} - \frac{1}{\Delta} \sum_j e^{\nu V_j} \frac{\partial G_j}{\partial \gamma_k}.$$

The derivative with respect to the homogeneity parameter μ is given by

$$\frac{\partial}{\partial \mu} \ln P = \frac{1}{G_i} \frac{\partial G_i}{\partial \mu} - \frac{1}{\Delta} \sum_j e^{\nu V_j} \frac{\partial G_j}{\partial \mu}$$

In general, the parameter μ is constraint to 1. However, Biogeme allows to estimate it if the user desires to do so. Therefore, the derivatives are necessary.

The derivative with respect to the scale parameter ν is given by

$$\frac{\partial}{\partial \nu} \ln P = V_i + \frac{1}{G_i} \frac{\partial G_i}{\partial \nu} - \frac{1}{\Delta} \frac{\partial \Delta}{\partial \nu}$$

where

$$\frac{\partial G_i}{\partial \nu} = \sum_j V_j e^{\nu V_j} \frac{\partial G_i}{\partial x_j}$$

and

$$\frac{\partial \Delta}{\partial \nu} = \sum_j \left(V_j e^{\nu V_j} G_j + e^{\nu V_j} \frac{\partial G_j}{\partial \nu} \right).$$

The GEV generating function for the MNL model is given by (14). We have

$$\frac{\partial G}{\partial y_i} = G_i = \mu y_i^{\mu-1},$$

and

$$\frac{\partial G}{\partial \mu} = \sum_{i=1}^n y_i^\mu \ln(y_i).$$

The second derivatives are

$$\frac{\partial^2 G}{\partial y_i \partial y_j} = \begin{cases} \mu(\mu - 1)y_i^{\mu-2} & \text{if } i = j \\ 0 & \text{if } i \neq j. \end{cases}$$

and

$$\frac{\partial^2 G}{\partial x_i \partial \mu} = (\mu \ln x_i + 1)x_i^{\mu-1}.$$

The GEV generating function for the NL model is given by (15). We have

$$\frac{\partial G}{\partial y_i} = \mu y_i^{\mu_{m_i}-1} \left(\sum_{j \in C_{m_i}} y_j^{\mu_{m_i}} \right)^{\left(\frac{\mu}{\mu_{m_i}} - 1\right)}$$

where m_i is the (unique) nest containing alternative i , and

$$\frac{\partial G}{\partial \mu} = \sum_{m=1}^M \frac{1}{\mu_m} \left(\sum_{i \in C_m} y_i^{\mu_m} \right)^{\frac{\mu}{\mu_m}} \ln \left(\sum_{i \in C_m} y_i^{\mu_m} \right).$$

The partial derivative with respect to one structural parameter μ_m is

$$\frac{\partial G}{\partial \mu_m} = \frac{\mu}{\mu_m} \left(\sum_{i \in C_m} y_i^{\mu_m} \right)^{\frac{\mu}{\mu_m}-1} \left(\sum_{i \in C_m} y_i^{\mu_m} \ln(y_i) \right) - \frac{\mu}{\mu_m^2} \left(\sum_{i \in C_m} y_i^{\mu_m} \right)^{\frac{\mu}{\mu_m}} \ln \left(\sum_{i \in C_m} y_i^{\mu_m} \right).$$

We have now the second partial derivative with respect to two variables i and j .

If $i = j$, we have

$$\frac{\partial^2 G}{\partial y_i^2} = \frac{\partial G_i}{\partial y_i} = \mu(\mu_m - 1)y_i^{(\mu_m-2)} \left(\sum_{i \in C_m} y_i^{\mu_m} \right)^{\left(\frac{\mu}{\mu_m} - 1\right)} + \mu(\mu - \mu_m)y_i^{(2\mu_m-2)} \left(\sum_{i \in C_m} y_i^{\mu_m} \right)^{\left(\frac{\mu}{\mu_m} - 2\right)}$$

If $i \neq j$ and $i, j \in C_m$, we have

$$\frac{\partial^2 G}{\partial y_i \partial y_j} = \frac{\partial G_i}{\partial y_j} = \mu(\mu - \mu_m)y_i^{\mu_m-1}y_j^{\mu_m-1} \left(\sum_{i \in C_m} y_i^{\mu_m} \right)^{\left(\frac{\mu}{\mu_m} - 2\right)}$$

If $i \in C_m$ and $j \notin C_m$, we have

$$\frac{\partial^2 G}{\partial y_i \partial y_j} = \frac{\partial G_i}{\partial y_j} = 0$$

The second partial derivative with respect to one variable and μ is given by

$$\frac{\partial^2 G}{\partial y_i \partial \mu} = \frac{\partial G_i}{\partial \mu} = y_i^{\mu_m - 1} \left(\sum_{i \in C_m} y_i^{\mu_m} \right)^{\frac{\mu}{\mu_m} - 1} \left(1 + \frac{\mu}{\mu_m} \ln \left(\sum_{i \in C_m} y_i^{\mu_m} \right) \right)$$

We have now the second partial derivative with respect to one structural parameter μ_m and one variable y_i . If $i \in C_m$, we have

$$\frac{\partial^2 G}{\partial y_i \partial \mu_m} = \frac{\partial G_i}{\partial \mu_m} = \mu \Gamma_m^{\frac{\mu}{\mu_m} - 1} y_i^{\mu_m - 1} \ln y_i + \mu y_i^{\mu_m - 1} \Gamma_m^{\frac{\mu}{\mu_m} - 1} \left(\frac{\mu}{\mu_m} - 1 \sum_j y_j^{\mu_m} \ln y_j - \frac{\mu}{\mu_m^2} \ln \Gamma_m \right),$$

where

$$\Gamma_m = \sum_{j \in C_m} y_j^{\mu_m}.$$

If $i \notin C_m$ we have

$$\frac{\partial^2 G}{\partial y_i \partial \mu_m} = \frac{\partial G_i}{\partial \mu_m} = 0.$$

The GEV generating function for the CNL model is given by (16). We have

$$G_i = \frac{\partial G}{\partial y_i} = \mu \sum_m \alpha_{im} y_i^{\mu_m - 1} \left(\sum_j \alpha_{jm} y_j^{\mu_m} \right)^{\frac{\mu}{\mu_m} - 1}.$$

The partial derivative with respect to μ , the homogeneity factor is

$$\frac{\partial G}{\partial \mu} = \sum_m \frac{1}{\mu_m} \Gamma_m^{\frac{\mu}{\mu_m}} \ln(\Gamma_m)$$

where

$$\Gamma_m = \sum_{j \in C_m} \alpha_{jm} y_j^{\mu_m}. \quad (24)$$

The partial derivative with respect to one nest parameter μ_m is

$$\frac{\partial G}{\partial \mu_m} = \frac{\mu}{\mu_m} \Gamma_m^{\frac{\mu}{\mu_m} - 1} \left(\sum_{j \in C_m} \alpha_{jm} y_j^{\mu_m} \ln(y_j) \right) - \frac{\mu}{\mu_m^2} \Gamma_m^{\frac{\mu}{\mu_m}} \ln(\Gamma_m)$$

and with respect α_{im} is

$$\frac{\partial G}{\partial \alpha_{im}} = \frac{\mu}{\mu_m} \Gamma_m^{\frac{\mu}{\mu_m} - 1} y_i^{\mu_m}$$

where Γ_m is defined by (24). We write now the second partial derivative with respect to two variables y_i and y_j . If $i = j$, we have

$$\frac{\partial^2 G}{\partial y_i^2} = \frac{\partial G_i}{\partial y_i} = \sum_m \frac{\mu}{\mu_m} \Gamma_m^{\frac{\mu}{\mu_m} - 2} \alpha_{im} \mu_m y_i^{\mu_m - 2} \left(\left(\frac{\mu}{\mu_m} - 1 \right) \alpha_{im} \mu_m y_i^{\mu_m} + \Gamma_m (\mu_m - 1) \right)$$

and if $i \neq j$, we have

$$\frac{\partial^2 G}{\partial y_i \partial y_j} = \frac{\partial G_i}{\partial y_j} = \sum_m \mu_m \mu \left(\frac{\mu}{\mu_m} - 1 \right) \alpha_{im} \alpha_{jm} \Gamma_m^{\frac{\mu}{\mu_m} - 2} y_i^{\mu_m - 1} y_j^{\mu_m - 1}$$

where

$$\Gamma_m = \sum_{j \in \mathcal{C}_m} \alpha_{jm} y_j^{\mu_m}.$$

The second partial derivative with respect to one variable y_i and μ is

$$\frac{\partial^2 G}{\partial y_i \partial \mu} = \frac{\partial G_i}{\partial \mu} = \sum_m \Gamma_m^{\frac{\mu}{\mu_m} - 1} \alpha_{im} y_i^{\mu_m - 1} \left(1 + \frac{\mu}{\mu_m} \ln(\Gamma_m) \right)$$

where Γ_m is defined by (24). The second partial derivative with respect to one nest parameter μ_m and one variable y_i is

$$\begin{aligned} \frac{\partial^2 G}{\partial y_i \partial \mu_m} = \frac{\partial G_i}{\partial \mu_m} = & - \frac{\mu}{\mu_m} \Gamma_m^{\frac{\mu}{\mu_m} - 1} \alpha_{im} y_i^{\mu_m - 1} \\ & - \frac{\mu^2}{\mu_m^2} \Gamma_m^{\frac{\mu}{\mu_m} - 1} \ln(\Gamma_m) \alpha_{im} y_i^{\mu_m - 1} \\ & + \frac{\mu}{\mu_m} \Gamma_m^{\frac{\mu}{\mu_m} - 1} \alpha_{im} y_i^{\mu_m - 1} \\ & + \mu \Gamma_m^{\frac{\mu}{\mu_m} - 1} \alpha_{im} y_i^{\mu_m - 1} \ln(y_i), \end{aligned}$$

and the second partial derivative with respect to α_{jk} and one variable y_i is

$$\frac{\partial^2 G}{\partial y_i \partial \alpha_{ik}} = \mu y_i^{\mu_k - 1} \Gamma_k^{\frac{\mu}{\mu_k} - 1} \left(1 + \alpha_{ik} \left(\frac{\mu}{\mu_k} - 1 \right) \Gamma_k^{-1} y_i^{\mu_k} \right)$$

and, if $i \neq j$,

$$\frac{\partial^2 G}{\partial y_i \partial \alpha_{jk}} = \mu \alpha_{ik} y_i^{\mu_k - 1} \left(\frac{\mu}{\mu_k} - 1 \right) \Gamma_k^{\frac{\mu}{\mu_k} - 2} y_j^{\mu_k}$$

where Γ_m is defined by (24).

Finally, we provide the derivatives of (9) with respect to the parameters. As the scale parameter ν has already been addressed above, we consider here

$$V(\beta, \lambda, \alpha) = \beta \frac{(x + \alpha)^\lambda - 1}{\lambda}$$

where the indices have been dropped for clarity. We have also

$$\lim_{\lambda \rightarrow 0} V(\beta, \lambda, \alpha) = \beta \ln(x + \alpha).$$

The derivatives are

$$\frac{\partial V}{\partial \beta} = \frac{(x + \alpha)^\lambda - 1}{\lambda},$$

$$\frac{\partial V}{\partial \lambda} = \frac{\beta}{\lambda^2} (1 + (x + \alpha)^\lambda (\lambda \ln(x + \alpha) - 1)),$$

$$\lim_{\lambda \rightarrow 0} \frac{\partial V}{\partial \lambda} = \frac{\beta}{2} \ln^2(x + \alpha)$$

and

$$\frac{\partial V}{\partial \alpha} = \beta(x + \alpha)^{\lambda-1}.$$

References

- Anderson, S. P., de Palma, A. and Thisse, J.-F. (1992). *Discrete Choice Theory of Product Differentiation*, MIT Press, Cambridge, Ma.
- Ben-Akiva, M. and Bierlaire, M. (1999). Discrete choice methods and their applications to short-term travel decisions, *in* R. Hall (ed.), *Handbook of Transportation Science*, Kluwer, pp. 5–34.
- Ben-Akiva, M. and Bolduc, D. (1996). Multinomial probit with a logit kernel and a general parametric specification of the covariance structure. Paper presented at the 3rd Invitational Choice Symposium, Columbia University.
- Ben-Akiva, M., Bolduc, D. and Walker, J. (2001). Specification, identification and estimation of the logit kernel (or continuous mixed logit) model. Working paper.
- Ben-Akiva, M., Bradley, M., Morikawa, T., Benjamin, J., Novak, T., Oppewal, H. and Rao, V. (1994). Combining revealed and stated preferences data, *Marketing letters* **5**(4): 335–350.
- Ben-Akiva, M. E. and Lerman, S. R. (1985). *Discrete Choice Analysis: Theory and Application to Travel Demand*, MIT Press, Cambridge, Ma.
- Ben-Akiva, M., McFadden, D., Train, K., Walker, J., Bhat, C., Bierlaire, M., Bolduc, D., Boersch-Supan, A., Brownstone, D., Bunch, D., Daly, A., de Palma, A., Gopinath, D., Karlstrom, A. and Munizaga, M. A. (2002). Hybrid choice models: Progress and challenges, *Marketing Letters* . accepted for publication.

- Ben-Akiva, M. and Morikawa, T. (1990). Estimation of travel demand models from multiple data sources, *Transportation and Traffic Theory* .
- Bhat, C. (2001). Quasi-random maximum simulated likelihood estimation of the mixed multinomial logit model, *Transportation Research B* **35**(7): 677–693.
- Bierlaire, M. (1998). Discrete choice models, in M. Labbé, G. Laporte, K. Tanczos and P. Toint (eds), *Operations Research and Decision Aid Methodologies in Traffic and Transportation Management*, Vol. 166 of *NATO ASI Series, Series F: Computer and Systems Sciences*, Springer Verlag, pp. 203–227.
- Bierlaire, M. (2001). A general formulation of the cross-nested logit model, *Proceedings of the 1st Swiss Transportation Research Conference*, Ascona, Switzerland.
- Bierlaire, M. (2002). The network GEV model, *Proceedings of the 2nd Swiss Transportation Research Conference*, Ascona, Switzerland.
- Bierlaire, M. (2003). An introduction to BIOGEME Version 0.6. <http://roso.epfl.ch/biogeme>.
- Daly, A. (2001). Recursive Nested EV model, *ITS Working Paper 559*, Institute for Transport Studies, University of Leeds.
- Gumbel, E. J. (1958). *Statistics of Extremes*, Columbia University Press, New York.
- Hensher, D. A. and Johnson, L. W. (1981). *Applied discrete choice modelling*, Croom Helm, London.
- Horowitz, J. L., Koppelman, F. S. and Lerman, S. R. (1986). *A self-instructing course in disaggregate mode choice modeling*, Technology Sharing Program, US Department of Transportation, Washington, D.C. 20590.
- Kuntsevich, A. V. and Kappel, F. (1997). *The Solver for Local Nonlinear Optimization Problems*, version 1.1 edn, Institute for Mathematics, Karl-Franzens University of Graz, Heinrichstr. 36, A-8010 Graz (Austria).
- Lawrence, C., Zhou, J. and Tits, A. (1997). User’s guide for CFSQP version 2.5: A C code for solving (large scale) constrained nonlinear (minimax) optimization problems, generating iterates satisfying all inequality constraints,

- Technical Report TR-94-16r1*, Institute for Systems Research, University of Maryland, College Park, MD 20742, 1997.
- Luce, R. (1959). *Individual choice behavior: a theoretical analysis*, J. Wiley and Sons, New York.
- McFadden, D. (1978). Modelling the choice of residential location, *in* A. Karlquist *et al.* (ed.), *Spatial interaction theory and residential location*, North-Holland, Amsterdam, pp. 75–96.
- McFadden, D. (1981). Econometric models of probabilistic choice, *in* C. Manski and D. McFadden (eds), *Structural analysis of discrete data with econometric application*, MIT Press, Cambridge, Mass.
- McFadden, D. and Train, K. (1997). Mixed multinomial logit models for discrete response, *Technical report*, University of California, Berkeley, Ca.
- Papola, A. (2000). Some development of the cross-nested logit model, *Proceedings of the 9th IATBR Conference*.
- Ramming, M. S. (2001). *Network Knowledge and Route Choice*, PhD thesis, Massachusetts Institute of Technology.
- Shor, N. (1985). *Minimization Methods for Non-Differentiable Functions*, Springer-Verlag.
- Small, K. (1987). A discrete choice model for ordered alternatives, *Econometrica* **55**(2): 409–424.
- Spellucci, P. (1998a). A new technique for inconsistent problems in the SQP method, *Math. Meth. of Oper. Res.* **47**: 355–400.
- Spellucci, P. (1998b). An SQP method for general nonlinear programs using only equality constrained subproblems, *MP* **82**: 413–448.
- Spellucci, P. (n.d.). *DONLP2 Users Guide*, Dpt of Mathematics, Technical University at Darmstadt, 64289 Darmstadt, Germany.
- Swait, J. (2001). Choice set generation within the generalized extreme value family of discrete choice models, *Transportation Research B* **35**(7): 643–666.

- Vovsha, P. (1997). Cross-nested logit model: an application to mode choice in the Tel-Aviv metropolitan area, Transportation Research Board, 76th Annual Meeting, Washington DC. Paper #970387.
- Vovsha, P. and Bekhor, S. (1998). The link-nested logit model of route choice: overcoming the route overlapping problem, *Transportation Research Record* **1645**: 133–142.
- Walker, J. L. (2001). *Extended discrete choice models: integrated framework, flexible error structures, and latent variables*, PhD thesis, Massachusetts Institute of Technology.
- Wen, C.-H. and Koppelman, F. S. (2001). The generalized nested logit model, *Transportation Research B* **35**(7): 627–641.

Example	\mathcal{L}^*	Time CFSQP	Time DONLP2	Time SOLVOPT
EX00	-1582.56	00:00:01	00:00:03	00:00:08
NGEV00	-1582.56	00:00:20	00:00:25	00:01:58
EX01	-1582.56	00:00:01	00:00:02	00:00:08
NGEV01	-1582.56	00:00:20	00:00:27	00:01:54
EX02	-1582.56	00:00:02	00:00:02	00:00:09
NGEV02	-1582.56	00:00:20	00:00:27	00:02:03
EX03	-1578.25	00:00:02	00:00:02	00:00:15
NGEV03	-1578.25	00:00:22	00:00:26	00:03:31
EX04	-1587.30	00:00:02	00:00:02	00:00:18
NGEV04	-1587.30	00:00:22	00:00:26	00:03:59
EX05	-1586.09	00:00:03	00:00:03	00:00:09
NGEV05	-1586.09	00:00:57	00:01:07	00:03:30
EX06	-691.937	00:00:05	00:00:06	00:00:16
NGEV06	-691.937	00:01:22	00:01:39	00:03:37
EX07	-690.833	00:00:07	00:00:11	00:00:20
NGEV07	-690.833	00:01:29	00:03:21	00:03:51
EX08	-688.665	00:00:07	00:00:08	00:00:48
NGEV08	-688.665	00:01:23	00:03:30	00:10:55
EX09	-691.21	00:00:10	00:00:13	00:01:07
NGEV09	-691.21	00:02:18	00:03:08	00:11:57
EX10	-658.205	00:23:42	00:03:23	****
NGEV10	-658.205	****	00:10:21	
EX11	-691.935	00:00:53	00:01:05	00:15:23
NGEV11	-691.935	00:09:07	00:14:43	
EX12	-662.619	00:00:10	00:01:36	00:30:22
NGEV12	-662.619	00:05:03	00:07:36	
EX13	-652.219	00:02:55	00:05:03	00:09:19
EX14	-676.072	00:03:25	00:05:49	00:11:53
EX15	-655.517	00:04:26	00:06:32	00:12:47
EX16	-657.651	00:05:00	00:08:43	00:13:46
EX17	-655.519	00:03:39	00:08:47	00:10:03
EX18	-685.451	00:04:55	00:07:30	00:14:15

Table 1: Comparison of the examples